

Neutrino scattering on polarized electron target and neutrino magnetic moment

Timur I. Rashba

E-mail: rashba@izmiran.rssi.ru

The Institute of the Terrestrial Magnetism, the Ionosphere and Radio Wave Propagation of the Russian Academy of Sciences IZMIRAN, Troitsk, Moscow region, 142190, Russia

Abstract

The completed and proposed experiments for the measurement of the neutrino magnetic moment are discussed. To improve the sensitivity of the search for the neutrino magnetic moment we suggest to use a polarized electron target in the processes of neutrino (antineutrino) – electron scattering. It is shown that in this case the weak interaction term in the total cross section is few times smaller comparing with unpolarized case, but the electromagnetic term does not depend on electron polarization.

The search of new bounds on the neutrino magnetic moment μ_ν in laboratory experiments would be very important for checking of new Physics beyond the Standard Model (SM) and for different astrophysical implications. The Resonant Spin-Flavor Precession (RSFP) scenario (Akhmedov, 1988; Lim and Marciano, 1988) in the case of a non-vanishing neutrino magnetic moment $\mu_\nu \neq 0$ is still considered as a possible solution to the Solar Neutrino Problem (SNP) (Berezinsky, 1998). The observations of neutron stars gave an information about the existence of strong magnetic fields (even more than the critical value $B_c = m_e^2/e = 4.41 \cdot 10^{13}$ G) which can interact with neutrino through its magnetic moment. The nature of a neutrino (Dirac or Majorana) determines the properties of neutrino magnetic moments, so a Dirac neutrino can have diagonal and transition (off-diagonal) moments, while Majorana neutrinos can have transition moments only (Schechter and Valle, 1981). All these hints claim direct laboratory measurements of the neutrino magnetic moment.

There is the laboratory bound on the electron neutrino magnetic moment from the experiments with reactor's neutrino combined by Derbin (1994), $\mu_{\nu_e} < 1.8 \cdot 10^{-10} \mu_B$. There are also different astrophysical constraints that are even more stringent then previous one. For the detailed discussion and references see Raffelt (1996).

To probe neutrino magnetic moment at a more lower level $\mu_{\nu_e} \sim 10^{-11} \mu_B$ some artificial neutrino sources of well-known activity and spectra can be used. There are some experiments planned with different possible artificial sources, e.g. ^{51}Cr (Ferrari et al., 1996), ^{90}Sr (Mikaelyan et al., 1998; Ianni and Montanino, 1999), ^{147}Pm (Kornoukhov, 1997; Barabanov et al., 1997), ^{55}Fe (Golubchikov et al., 1996). The recently proposed measurement of neutrino magnetic moment at a level better than $10^{-11} \mu_B$ ($\sim 3 \cdot 10^{-12} \mu_B$) with the use of the tritium neutrino emitter (antineutrino source with the energy spectrum endpoint 18.6 keV) and a semiconductor cryodetector is planned to reach extra low threshold about 1.1 eV (Trofimov et al., 1998; Bogdanova, 2000) (see Table 1).

Reference	$\nu(\bar{\nu})$ source	Detector	$\mu_\nu/\mu_B >$
Ferrari et al., 1996	^{51}Cr	BOREXINO	$5 \cdot 10^{-11}$
Ianni et al., 1999	$^{51}\text{Cr}, ^{90}\text{Sr}$	BOREXINO	$6 \cdot 10^{-11}$
Miranda et al., 1999	$^{51}\text{Cr}, ^{90}\text{Sr}$	HELLAZ	$6 \cdot 10^{-11}$
Beda et al., 1998	Reactor	Ge-NaI	$3 \cdot 10^{-11}$
TEXONO, Wong, 1999	Reactor, Taiwan	CsI	$3 \cdot 10^{-11}$
Kurchatov-PNPI, Kozlov, 1999	Krasnoyarsk reactor	Si	$2 \cdot 10^{-11}$
MUNU, Brogini, 1999	Bugey reactor	CF_4	$2 \cdot 10^{-11}$
Mikaelyan et al., 1998	^{90}Sr	BOREXINO	$1.5 \cdot 10^{-11}$
Golubchikov et al., 1996	^{55}Fe	Si	$5 \cdot 10^{-12}$
Kornoukhov, 1997	^{147}Pm	Ge, Si or NaI	$5 \cdot 10^{-12}$
Barabanov et al., 1997	^{147}Pm	NaI	$3 \cdot 10^{-12}$
Trofimov et al., 1998	TiT_2	Si	$3 \cdot 10^{-12}$

Table 1. Proposed and planning experiments and neutrino magnetic moments which can be measured.

We suggest to use a polarized electron target for more precise measurements of a neutrino magnetic moment in the processes of neutrino (antineutrino) – electron scattering (Rashba and Semikoz, 2000). Kinematics of such scattering for which the recoil electron energy $T_e = E_2 - m_e$ should be measured is shown in Fig.1. We have fixed $z0x$ -plane based on the initial neutrino momentum $\vec{k}_1$ directed along z -axis and on the 3-vector part $\vec{\xi}_e$ of the four-spin $a_\mu = (0, \vec{\xi}_e)$ at the rest frame of the initial electron ($\vec{p}_1 = 0$) entering the Mishel–Wightman density matrix, $\rho(p_1) = (\hat{p}_1 + m_e)(1 + \gamma_5 \hat{a})$.

The cross section of the electron neutrino (antineutrino) scattering off electrons consists of two terms: weak and electromagnetic ones without interference term that can be neglected in the massless limit $m_\nu \rightarrow 0$,

$$\left(\frac{d\sigma}{dT_e d\phi} \right) = \left(\frac{d\sigma}{dT_e d\phi} \right)^{\text{weak}} + \left(\frac{d\sigma}{dT_e d\phi} \right)^{\text{em}}. \quad (1)$$

After integration over the azimuthal angle ϕ (see Fig.1) the weak cross sections have the following forms:

$$\begin{aligned} \left(\frac{d\sigma}{dT_e} \right)_\nu^{\text{weak}} &= \frac{2G_F^2 m_e}{\pi} \left[g_{eL}^2 \left(1 + |\vec{\xi}_e| \cos \theta_\xi \right) + g_R^2 \left(1 - \frac{T_e}{\omega_1} \right)^2 \times \right. \\ &\times \left. \left(1 - |\vec{\xi}_e| \cos \theta_\xi \left(1 - \frac{m_e T_e}{\omega_1 (\omega_1 - T_e)} \right) \right) - g_{eL} g_R \frac{m_e T_e}{\omega_1^2} \left(1 + |\vec{\xi}_e| \cos \theta_\xi \right) \right] \end{aligned} \quad (2)$$

for left-handed neutrino and

$$\begin{aligned} \left(\frac{d\sigma}{dT_e} \right)_{\bar{\nu}}^{\text{weak}} &= \frac{2G_F^2 m_e}{\pi} \left[g_R^2 \left(1 - |\vec{\xi}_e| \cos \theta_\xi \right) + g_{eL}^2 \left(1 - \frac{T_e}{\omega_1} \right)^2 \times \right. \\ &\times \left. \left(1 + |\vec{\xi}_e| \cos \theta_\xi \left(1 - \frac{m_e T_e}{\omega_1 (\omega_1 - T_e)} \right) \right) - g_{eL} g_R \frac{m_e T_e}{\omega_1^2} \left(1 - |\vec{\xi}_e| \cos \theta_\xi \right) \right] \end{aligned} \quad (3)$$

for right-handed antineutrino, where $|\vec{\xi}_e|$ is the value of the initial electron polarization. Here G_F is the Fermi constant; $g_{eL} = \sin^2 \theta_W + 0.5$, $g_R = \sin^2 \theta_W$ ($\sin^2 \theta_W \approx 0.23$) are the couplings in the SM; $\omega_1 = k_1$ is the initial neutrino energy; θ_ξ is the angle between the neutrino momentum $\vec{k}_1$ and the electron polarization vector $\vec{\xi}_e$. If the new (experimental) parameter $|\vec{\xi}_e| \cos \theta_\xi$ vanishes, $|\vec{\xi}_e| \cos \theta_\xi \rightarrow 0$, our results convert to the standard ones (Okun', 1990).

Electromagnetic terms of the neutrino (antineutrino) scattering cross section do not depend on the electron polarization $\vec{\xi}_e$ and have the same form for neutrinos and antineutrinos. If CP holds¹ the electromagnetic

¹In the Dirac neutrino case when CP conserves the diagonal electric dipole moment is absent, $d_\nu = 0$. For Majorana neutrino this form is valid if CP parities of initial and final neutrino states are opposite hence the electric dipole moment does not contribute, $d_{ij} = 0$ (Schechter and Valle, 1981). For the Dirac neutrino transition moment we should substitute $|\mu_\nu|^2 \rightarrow |\mu_{ij} - id_{ij}|^2$ where μ and d are not separated and obey the equality $\text{Re } \mu \text{ Re } d + \text{Im } \mu \text{ Im } d = 0$ (Raffelt, 1996).

term in Eq. (1) can be written as

$$\left(\frac{d\sigma}{dT_e}\right)^{\text{em}} = \frac{\pi\alpha^2}{m_e^2} \left(\frac{1}{T_e} - \frac{1}{\omega_1}\right) \frac{|\mu_\nu|^2}{\mu_B^2}, \quad (4)$$

here μ_B is the Bohr magneton.

As well as for the case of an unpolarized target ($\vec{\xi}_e = 0$) the electromagnetic interaction via a large magnetic moment increases very significantly in the region of small energy transfer $T_e = \omega_1 - \omega_2 \ll m_e$, ω_1 and can be comparable or greater than the weak interaction. This gives a possibility to determine an upper limit on the neutrino magnetic moment or to measure it.

Weak interaction cross sections (Eqs. (2) and (3)) occur very sensitive both to the polarization of electron target $|\vec{\xi}_e|$ and to the angle θ_ξ between the neutrino momentum and the initial electron polarization. For the known values of the SM parameters ($g_{eL}^2 \approx 0.533$, $g_R^2 \approx 0.053$ and $g_{eL}g_R \approx 0.168$) one can observe from Eqs. (2) and (3) a new possibility for decreasing of the main weak term ($\sim g_{eL}^2$) in the cross section Eq. (1) choosing large polarization values ($|\vec{\xi}_e| \rightarrow 1$) and with a choice of the specific geometry ($\theta_\xi = \pi$). The weak cross sections for ^{51}Cr neutrino source and different values of the initial electron polarization are shown in Fig.3.

In the case discussed above ($|\vec{\xi}_e| \cos \theta_\xi = -1$) the weak interaction cross sections are proportional to g_R^2 . It means that target electrons (in laboratory system) that are fully polarized in the opposite direction to the neutrino momentum will interact with a neutrino as right chiral particles. This can be explained by the following way. Let's make the Lorentz boost from the laboratory system to the ultrarelativistic one which moves along neutrino momentum with $V \sim c = 1$. In this system initial electron is an ultrarelativistic particle and it can be considered as a massless one. Since the modulo of electron polarization is Lorentz-invariant, $|\vec{\xi}_e| = \sqrt{-a_\mu a^\mu}$, and the electron momentum direction appeared in chosen system is parallel to its polarization vector such electron is the right-handed particle (right-helicity = +1) having same chirality (+1) in massless limit.

The best case of the antiparallel neutrino (antineutrino) beam $\cos \theta_\xi = -1$ can be generalized for a more realistic case with an integration over θ_ξ accounting for a concrete geometry of the artificial (isotope) neutrino source placed outside of the detector with known sizes.

Note that the polarization contribution to the $\tilde{\nu}e$ -scattering vanishes at the two points:

$$T_{e1} = \frac{\omega_1^2}{m_e + \omega_1} \left(1 + \frac{g_R}{g_{eL}}\right), \quad T_{e2} = \omega_1 \left(1 - \frac{g_R}{g_L}\right). \quad (5)$$

For the tritium antineutrino source the polarization reduces (enhances) the unpolarized part of the SM weak cross section below (above) T_{e1} . For the same source the second root T_{e2} occurs out of the kinematic allowed region, $T_{e2} > T_{\text{max}} = 2\omega_1^2/(m_e + 2\omega_1)$. Hence in the case of the $\tilde{\nu}e$ -scattering the low energy region $T_e < T_{e1}$ is preferable to look for an electromagnetic effect from the sum Eq. (1).

One can easily check that in the case of the νe -scattering ($g_{eL} \leftrightarrow g_R$) both roots Eq. (5) are out of the kinematic allowed region and the polarization term reduces the SM weak cross-section for any energy T_e .

Let us consider the spectra of recoil electrons from an antineutrino (neutrino) source calculated via the averaging of the differential cross-sections over the antineutrino (neutrino) spectrum $\rho(\omega_1)$,

$$\begin{aligned} S_{\text{free}}^{\text{weak}}(T_e) &= \int_{\omega_{1\text{min}}} d\omega_1 \frac{d\sigma^{\text{weak}}(T_e, \omega_1)}{dT_e} \rho(\omega_1), \\ S_{\text{free}}^{\text{em}}(T_e) &= \int_{\omega_{1\text{min}}} d\omega_1 \frac{d\sigma^{\text{em}}(T_e, \omega_1)}{dT_e} \rho(\omega_1), \end{aligned} \quad (6)$$

where weak and electromagnetic cross-sections are given by Eqs. (2), (3) and Eq. (4) correspondingly and $\omega_{1\text{min}} = (T_e/2)[1 + \sqrt{1 + 2m_e/T_e}]$ is the minimal neutrino energy given by the kinematic limit $T_e \leq T_{\text{max}}$. The ratio of total recoil electron spectra to weak one for different values of initial electron polarization and neutrino magnetic moment for the tritium antineutrino source is shown in Figures 2, 6 and 7. The same plots for the ^{51}Cr neutrino source are shown in Figures 4 and 5.

It was found by Kopeikin et al. (1997), that inelastic spectra $S_{\text{in}}(q)$ which depend on the energy transfer $q = \varepsilon_i + T_e$ for a bound initial electron knocked out from the i -shell coincide with the spectra Eq. (6), $S_{\text{in}}(q) = S_{\text{free}}(q)\theta(q - \varepsilon_i)$, when the value q is larger than the binding energy ε_i , $q > \varepsilon_i$. Really, q is exactly the event energy measured in the experiment since soft X-quanta and mainly Auger electrons emitted by an atom from which a recoil electron was knocked out are automatically absorbed in the detector fiducial volume or their total energy ε_i adds to the kinetic energy T_e (Kopeikin et al., 1997).

For shells ordered, $i = K, L, M, N, O, \dots$, let us consider low energy transfer, $\varepsilon_{i-1} > q \geq \varepsilon_i$, when antineutrinos (neutrinos) knock out outer (let us say $i = O$, main Bohr number $n = 5$) electrons only not touching all inner ones. Moreover, since inner electrons from filled shells with $J = L = S = 0$ ($i - 1 = K, L, \dots$) do not contribute to the electron polarization one has a sense to consider only the νe -scattering off outer i -electrons for which the Zeeman splitting energy is given by

$$E_{J, M_J} = g_J \mu_B M_J H, \quad (7)$$

where g_J is the Lande factor and $M_J = -J, \dots, J-1, J$ is the projection of the total spin $\vec{J}$ on the magnetic field direction.

In order to obey such conditions we propose to use a tritium antineutrino source ($T_{\text{max}} \approx 1.26$ keV) and a cryogenic detector with the lowest threshold ($\sim$ eV-region) for which some lower energy bins $T_{\text{th}} \leq T_{\text{th}} + \Delta T_e \leq T_{\text{th}} + 2\Delta T_e, \dots$ within the interval

$$T_{\text{th}} < q = T_e + \varepsilon_i < \varepsilon_{i-1}, \quad (8)$$

can be separated. For instance, such low thresholds are considered by Trofimov et al. (1998) for the Si-semiconductor detector without magnetic fields. Hence the recoil electron energy T_e should be lower than *the difference between the binding energy of the last filled shell ε_{i-1} and outer (valence) electron energy ε_i* , $T_e < \varepsilon_{i-1} - \varepsilon_i$. In such case only polarized (outer) electrons contribute to the event spectrum to be measured.

One assumes that a low bin width $\Delta T_e \sim T_{\text{th}} \sim$ a few eV would be enough to measure the spectrum $S_{\text{in}}(q) = S_{\text{free}}(q)$.

Note that we have discussed above the spectrum $S_{\text{in}}(q)$ for the ionization process $\nu + e(i) \rightarrow \nu + e$ for an individual electron on the i -subshell with a free final electron. Accounting for all polarized electrons on that outer i -subshell of an atom Z (ion in the detector molecule) we find that the total sum over subshells (Kopeikin, 1997) reduces within the low-energy interval Eq. (8) to the fraction of polarized electrons:

$$\frac{S_{\text{in}}(q)}{S_{\text{free}}(q)} = \frac{1}{Z} \sum_{i=K, L, M, N, O, \dots} n_i \theta(q - \varepsilon_i) \rightarrow \frac{n_O}{Z}, \quad (9)$$

where n_i is the number of electrons on the i -subshell, n_O is the number of polarized (valence) electrons on the last unfilled, let us say, $i = O$ -shell.

The equal reduction of weak and electromagnetic inelastic contributions $S_{\text{in}}(q)$ comparing with the hard energy case $q \gg \varepsilon_i$ (for which $S_{\text{in}} \approx S_{\text{free}}$) leads to a calculable event statistics decrease. Nevertheless, for low energies Eq. (8) near T_{th} the conserving ratio

$$\frac{S_{\text{in}}^{\text{em}}(q) + S_{\text{in}}^{\text{weak}}(q)}{S_{\text{in}}^{\text{weak}}(q)} = \frac{S_{\text{free}}^{\text{em}}(q) + S_{\text{free}}^{\text{weak}}(q)}{S_{\text{free}}^{\text{weak}}(q)} \quad (10)$$

given by Eq. (6) is still more sensitive to the neutrino magnetic moment in the case of polarized outer electrons than for unpolarized targets discussed in (Trofimov et al., 1998) (compare in Fig. 2).

The valence electrons can be fully polarized. Really, applying realistic magnetic fields $H \gtrsim 10^4$ G to a cryodetector one can easily reach the ratio $\mu_B H/T > 1$ at low temperatures $T \lesssim 0.5$ K. Known experimental methods (Groot et al., 1965) allow to reach high electron polarization at temperatures less than the Curie point T_C when the parameter $\mu_B H/T$ is quite large.

For such conditions one can expect 100% polarization of outer (valence) electrons in the sum over M_J entering the electron polarization $|\vec{\xi}_e| = P$,

$$P = \frac{\sum_{|M_J|} [\exp(g_J \mu_B |M_J| H/T) - \exp(-g_J \mu_B |M_J| H/T)]}{\sum_{|M_J|} [\exp(g_J \mu_B |M_J| H/T) + \exp(-g_J \mu_B |M_J| H/T)]}. \quad (11)$$

Here we have used the Boltzman distribution for independent ions (atoms) for which the fraction of the polarized electrons with the total spin projection M_J is given by the ratio $N_{M_J}/N_0 = \exp(-g_J\mu_B M_J H/T)$ where N_0 is the normalization factor not playing a role in the ratio Eq. (11).

Now let us turn to some experimental prospects. Such weakly interacting Boson gas as atomic hydrogen H^2 being spin-polarized in high magnetic fields is used for possible Bose-Einstein condensation at low temperatures (Silvera, 1995) and seems could be also used as an ideal polarized (100 %) electron target for the νe -scattering. However, we do not know how to register recoil electrons there.

The Neganov-Trofimov-Luke effect of ionization-to-heat conversion (Neganov and Trofimov, 1981; Luke, 1988) that was proposed for the use of cryogenic semiconductor detectors to measure lowest neutrino magnetic moments (Golubchikov et al., 1996; Trofimov et al., 1998) may be modified for such magnetic semiconductors as ferromagnets EuS, EuO where outer f -electrons are fully polarized along an external magnetic field (Nagaev, 1979).

The method should be similar to Neganov and Trofimov (1981) and Luke (1988) but has some distinctions. At the initial state and low temperatures $T < T_C$ ($T_C \sim 14$ K for EuS and $T_C \sim 69$ K for EuO) there are no electrons (holes) in conductive (valence) zones, $n_e = n_h \sim \exp(-E_g/T) \approx 0$, $E_g \gg T$. Then a neutrino hits a polarized valence electron which overcomes the semiconductor gap E_g ($E_g = 0.9$ eV for EuO or $E_g = 1.5$ eV for EuS) appearing in the conductive zone where it could be accelerated by an external electric field of the known value applied to the crystal to get a measurable energy ΔE_e . In the case of a simple semiconductor like Si kept at the thermostat temperature T the following electron-phonon interaction leads to the heating of such Si-crystal for which a low lattice heat capacity $c_v \sim (T/T_D)^3$ allows to get a large temperature jump $\Delta T \sim \Delta E_e (T_D/T)^3$ caused by the electron energy absorption ($\sim \Delta E_e$).

On the other hand, in a ferromagnet semiconductor the lattice heat capacity scales as $c_v \sim (T/T_C)^{3/2}$ (Nagaev, 1979; Ziman, 1972; White, 1970) where the Curie temperature T_C is much lower than the Debye one T_D , $T_C \ll T_D$. As result the electron-magnon interaction heats crystal (through spin-waves excited by a recoil electron) weaker than it happens through phonons in the Si-case. From this we conclude that new cryogenic detectors suggested here at least should be kept at lower temperatures while there are other possibilities that are considered now (e.g. ferromagnet semiconductor $CdCr_2Se_4$ with higher $T_C \sim 110$ K, etc).

Note that in the low energy transfer region Eq. (8) the weak cross sections in the case of maximal (100 %) electron polarization and for the opposite direction of the magnetic field $\vec{H}$ with respect to the initial neutrino (antineutrino) momentum ($|\vec{\xi}_e| \cos \theta_\xi = -1$) are 5 times smaller than weak cross sections for the unpolarized case ($|\vec{\xi}_e| = 0$). The ratios Eq. (10) of the total recoil electron spectra and the weak one for the tritium antineutrino scattering off polarized electrons are shown in Figures 6 and 7.

In a real experiment the parameter $|\vec{\xi}_e| \cos \theta_\xi$ may in practice not equal to -1 because of the partially polarized electrons $|\vec{\xi}_e| < 1$ as well as due to the non-collinear geometry of the experiment ($\theta_\xi \neq \pi$). E.g. the ratio of the total cross section and the weak one for $\theta_\xi \approx 155$ degrees are shown in Figures 4–7 where the polarization parameter $|\vec{\xi}_e| \cos \theta_\xi$ is about -0.9 for $|\vec{\xi}_e| = 1$.

The accuracy of $\mu\nu$ -calculations depends also on experimental uncertainties including a background, an isotope source activity, a concrete geometry of the experiment, etc. (see e.g. Ianni and Montanino (1999)) that could be done for a future concrete polarized cryodetector with a low threshold and is beyond the scope of the present proposal.

Note that we have omitted radiative corrections (RC) in the main term $g_{eL}^2(1 + \alpha f_-/\pi)(1 + |\vec{\xi}_e| \cos \theta_\xi)$ (Bahcall et al., 1995) entering Eq. (2) where the RC term $\alpha f_-/\pi$ is expected at the level ~ 1 % in the low energy region, or occurs at the negligible level comparing with the strong polarization influence the weak cross-section.

I am grateful to my co-author V.B.Semikoz for many discussions and help in this work. I thank M.I.Visotsky, V.A.Novikov and D.S.Gorbunov for useful discussions during the School. I am grateful to the Organizing Committee of the ITEP Winter School 2000 for the pleasant and interesting atmosphere. This work was supported by the RFBR grant 00-02-16271.

²Molecular hydrogen H_2 is diamagnetic.

References

- Akhmedov E.Kh., "Resonant amplification of neutrino spin rotation in matter and the solar neutrino problem", *Phys. Lett.*, **B 213**, 64 (1988).
- Bahcall J.N., Kamionkowski M. and Sirlin A., "Solar neutrinos: radiative corrections in neutrino - electron scattering experiments", *Phys. Rev.*, **D 51**, 6146 (1995); e-print Archive: astro-ph/9502003.
- Barabanov I.R. et al., *Astropart. Phys.*, **8**, 67 (1997).
- Beda A.G., Demidova E.V., Starostin A.S., Voloshin M.B., "On the feasibility of low-background Ge-NaI spectrometer for neutrino magnetic moment measurement", *Phys. Atom. Nucl.*, **61**, 66-73 (1998) [*Yad. Fiz.*, **61**, 72-79 (1998)].
- Berezinsky V., "Oscillation solutions to solar neutrino problem", Talk given at 19th Texas Symposium on Relativistic Astrophysics: Texas in Paris, France, 14-18 Dec 1998; e-Print Archive: hep-ph/9904259.
- Bogdanova L.N., "Studying of the neutrino properties with a superstrong tritium source", in the Proceedings (2000).
- Broggini, C., "Status of the MUNU experiment", *Nucl. Phys. Proc. Suppl.*, **70**, 188 (1999).
- Derbin A.V. et al., "Restriction on the magnetic dipole moment of reactor neutrinos", *Yad. Fiz.*, **57**, 236 (1994) [*Phys. Atom. Nucl.*, **57**, 222 (1994)].
- Ferrari N., Fiorentini G. and Ricci B., "The Cr-51 neutrino source and Borexino: a desirable marriage", *Phys. Lett.*, **B 387**, 427 (1996).
- Golubchikov A.V., Zaimidoroga O.A., Smirnov O.Y. and Sotnikov A.P., "On neutrino magnetic moment measurement using artificial neutrino source", *Phys. Atom. Nucl.*, **59**, 1916 (1996).
- de Groot S.R., Tolhoek H.A. and Huiskamp W.I. in *Alpha-, beta- and gamma-ray spectroscopy*, edited by Siegbahn K., Vol. 3, North-Holland Publishing Company, Amsterdam (1965).
- Ianni A. and Montanino D., "The Cr-51 and Sr-90 sources in BOREXINO as tool for neutrino magnetic moment searches", *Astropart. Phys.*, **10**, 331 (1999).
- Kopeikin V.I., Mikaelyan L.A., Sinev V.V. and Fayans S.A., "Reactor antineutrino-electron scattering", *Yad. Fiz.*, **60** 2032 (1997).
- Kornoukhov V.N., "Laboratory source of antineutrinos on the basis of Pm-147: conceptual design and applications", *Phys. Atom. Nucl.*, **60**, 558 (1997).
- Kozlov Yu.V. et al., "Today and future neutrino experiments at Krasnoyarsk nuclear reactor", e-Print Archive: hep-ex/9912046 (1999).
- Luke P., *J. Appl. Phys.*, **64**, 6858 (1988).
- Lim C.-S. and Marciano W.J., "Resonant spin - flavor precession of solar and supernova neutrinos", *Phys. Rev.*, **D 37**, 1368 (1988).
- Mikaelyan L.A., Sinev V.V. and Fayans S.A., "On a precise check of the standard model in an experiment with a Sr-90 beta source", *JETP Lett.*, **67**, 453 (1998).
- Miranda O.G., Segura J., Semikoz V.B., Valle J.W.F., "Probing neutrino magnetic moments at underground detectors with artificial neutrino sources", e-Print Archive: hep-ph/9906328.
- Nagaev E.L., *Fizika magnitnykh poluprovodnikov*, "Nauka", Moscow, (1979), in Russian [Nagaev E.L., *Physics of magnetic semiconductors*, translated by M. Samokhvalov, ed. by the author, Mir Publishers, Moscow, 388 pp. (1983)].

- Neganov B.S., Trofimov V.N., *URSS patent* 1037771 (1981), *Otkrytiya, Izobret.*, **146**, 215 (1985).
- Okun' L.B., *Leptons and quarks*, "Nauka", Moscow (1990).
- Raffelt G.G., *Stars as Laboratories for Fundamental Physics*, The University of Chicago Press (1996).
- Rashba T.I. and Semikoz V.B., "Neutrino scattering on polarized electron target as a test of neutrino magnetic moment", *Phys. Lett. B*, **479**, 218 (2000).
- Silvera I.F., *Journal of Low Temperature Physics*, **101**, 49 (1995).
- Trofimov V.N., Neganov B.S. and Yukhimchuk A.A., "Measurement of the neutrino magnetic moment at a level better than $10^{-11}\mu_B$ with a tritium $\bar{\nu}$ emitter and cryodetector (project)", *Yad. Fiz.*, **61** 1373 (1998) [*Phys. Atom. Nucl.*, **61** 1271 (1998)].
- Schechter J. and Valle J.W.F., "Majorana neutrinos and magnetic fields", *Phys. Rev.*, **D 24**, 1883 (1981); **D 25**, 283 (1982).
- White R.M., *Quantum Theory of Magnetism*, McGraw-Hill Book Company, New York (1970), formula (6.64).
- Wong H.T., Li J., "A pilot experiment with reactor neutrinos in Taiwan", *Nucl. Phys. B (Procs. Suppl.)*, **77**, 177 (1999).
- Ziman J.M., *Principles of the Theory of Solids*, Cambridge, The University Press, (1972), Chapter 10, § 11;

Figures

Figure 1. Kinematics of the neutrino scattering off the polarized electron.

Figure 2. The ratio of the total recoil electron spectrum and the weak one (Eq. 10) for different values of the polarization parameter $|\vec{\xi}_e| \cos \theta_\xi = 0, -1$ and for the fixed neutrino magnetic moment $\mu_\nu = 3 \cdot 10^{-12} \mu_B$ is shown as dashed and solid lines, correspondingly. Tritium antineutrino emitter.

Figure 3. The weak interaction term of the recoil electron spectrum one (Eq. 10) for different values of the polarization parameter $|\vec{\xi}_e| \cos \theta_\xi = 0, -0.9, -1$ is shown by dash-dotted, dashed and solid lines correspondingly. The dotted line shows the electromagnetic term of the recoil electron spectrum for the neutrino magnetic moment $\mu_\nu = 3 \cdot 10^{-12} \mu_B$. Neutrino source is ^{51}Cr .

Figure 4. The ratio of the total recoil electron spectrum and the weak one (Eq. 10) for different values of the polarization parameter $|\vec{\xi}_e| \cos \theta_\xi = 0, -0.9, -1$ and for the fixed neutrino magnetic moment $\mu_\nu = 10^{-12} \mu_B$ is shown by short-dashed, dashed and solid lines, correspondingly. The zero neutrino magnetic moment case is shown by dotted line. Neutrino source is ^{51}Cr .

Figure 5. The ratio of the total recoil electron spectrum and the weak one (Eq. 10) for fixed maximal value of polarization parameter $|\vec{\xi}_e| \cos \theta_\xi = -1$ and for different neutrino magnetic moments $\mu_\nu = 0, 3 \cdot 10^{-13} \mu_B, 10^{-12} \mu_B, 2 \cdot 10^{-12} \mu_B$ is shown by dotted, short-dashed, dashed and solid lines, correspondingly. Neutrino source is ^{51}Cr .

Figure 6. The ratio of the total recoil electron spectrum and the weak one (Eq. 10) for different values of the polarization parameter $|\vec{\xi}_e| \cos \theta_\xi = 0, -0.9, -1$ and for the fixed neutrino magnetic moment $\mu_\nu = 3 \cdot 10^{-13} \mu_B$ is shown by short-dashed, dashed and solid lines, correspondingly. The zero neutrino magnetic moment case is shown by dotted line. Tritium antineutrino emitter.

Figure 7. The ratio of the total recoil electron spectrum and the weak one (Eq. 10) for fixed maximal value of polarization parameter $|\vec{\xi}_e| \cos \theta_\xi = -1$ and for different neutrino magnetic moments $\mu_\nu = 0, 10^{-13} \mu_B, 3 \cdot 10^{-13} \mu_B, 10^{-12} \mu_B$ is shown by dotted, short-dashed, dashed and solid lines, correspondingly. Tritium antineutrino emitter.




Figure 1




Figure 2.




Figure 3.




Figure 4. Recoil electron kinetic energy T (eV)




Figure 5. Recoil electron kinetic energy T (eV)




Figure 6.




Figure 7.